



Whelton, H. L., Lewis, J., Halstead, P., Isaakidou, V., Triantaphyllou, S., Tzevelekidi, V., ... Evershed, R. P. (2018). Strontium isotope evidence for human mobility in the Neolithic of northern Greece. *Journal of Archaeological Science: Reports*, 20, 768-774.
<https://doi.org/10.1016/j.jasrep.2018.06.020>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.jasrep.2018.06.020](https://doi.org/10.1016/j.jasrep.2018.06.020)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.jasrep.2018.06.020> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Strontium isotope evidence for human mobility in the Neolithic of northern Greece

Whelton, H. L.^a, Lewis, J.^b, Halstead, P.^c, Isaakidou, V.^d, Triantaphyllou, S.^e, Tzevelekidi, V.^f,
Kotsakis, K.^e, Evershed, R. P.^a

^a*Organic Geochemistry Unit, School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8
ITS, UK*

^b*Bristol Isotope Group, School of Earth Sciences, University of Bristol, Wills Memorial Building,
Queen's Road, Bristol, BS8 1RJ, UK*

^c*Department of Archaeology, University of Sheffield, Sheffield, UK*

^d*School of Archaeology, University of Oxford, Oxford, UK*

^e*School of History & Archaeology, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece*

^f*Ephorate of Antiquities of West Attica, Pireus and Islands, Ministry of Culture and Sports, Athens,
Greece*

Abstract

Strontium isotope ratios are widely used in archaeology to differentiate between local and non-local populations. Herein, strontium isotope ratios of 36 human tooth enamels from seven archaeological sites spanning the Early to Late Neolithic of northern Greece (7th-5th millennia B.C.E.) were analysed with the aim of providing new information relating to the movement of humans across the region. Local bioavailable ⁸⁷Sr/⁸⁶Sr signals were established using tooth enamel from 26 domestic animals from the same Neolithic sites. ⁸⁷Sr/⁸⁶Sr values of faunal samples correlate well with predicted strontium isotope ratios of the local geology. This is consistent with animal management occurring at a local level, although at Late Neolithic sites strontium isotope values became more varied, potentially indicating changing herding practices. The strontium isotope analysis of human tooth enamel likewise suggests limited population movement within the Neolithic of northern Greece. Almost all individuals sampled exhibited ⁸⁷Sr/⁸⁶Sr values consistent with having spent their early life (during the period of tooth mineralisation) in the local area, although movement could have occurred between isotopically homogeneous areas. The strontium isotope ratios of only three individuals lay outside of the local bioavailable ⁸⁷Sr/⁸⁶Sr range and these individuals are interpreted as having spent their early lives in a region with a more radiogenic biologically available ⁸⁷Sr/⁸⁶Sr. Mobility patterns determined using Sr isotope analysis supports the current evidence for movement and exchange observed through studies of pottery circulation. Suggesting limited movement in the Early and Middle Neolithic and greater movement in the Late Neolithic.

Keywords

Strontium isotopes, Neolithic, Greece, Mobility, Human, Animals

Highlights

- Sr isotope ratios were determined for 36 human and 26 faunal tooth enamel samples
- Seven sites spanning the EN to the LN of northern Greece
- $^{87}\text{Sr}/^{86}\text{Sr}$ values of human tooth enamel shows that population movement was limited
- Three individuals identified as ‘non-locals’
- Sr ratio of faunal enamel infers animal management occurred on a local level

1. Introduction

The application of strontium isotope analyses to archaeological skeletal remains can provide information regarding the movement of humans and animals by comparing the strontium isotope signature of an individual to the biologically available signature determined by the surrounding biosphere (Bentley, 2006). Given suitable variation in the biologically available signatures in a region, strontium isotope ratios are able to differentiate between local and non-local populations. This can be used as complementary evidence to that obtained from the traded and exchanged goods, identified through material cultural remains, which together can provide a record of movement and exchange networks across a region. Strontium isotope analyses have thus been widely utilised to explore the mobility of early Hominins (Balter *et al.*, 2012), Pliocene mammals (Hoppe *et al.*, 1999), Neanderthal populations (Britton *et al.*, 2011) and Pre-historic human and animal mobility (Grupe *et al.*, 1997; Viner *et al.*, 2010; Bentley, 2013; Boric and Price, 2013; Giblin *et al.*, 2013; Gerling, 2015; Henton *et al.*, 2017).

Evidence for movement and exchange has been identified in the Greek Neolithic from the movement of material culture, including pottery, stone tools and shell ornaments. The study of fine-decorated pottery assemblages has shown that the circulation of pottery in northern Greece was very limited (Pentedeke, 2011; Urem-Kotsou *et al.*, 2012). Due to the low amount of pottery found in the Early Neolithic (EN) it is difficult to ascertain if exchanges of pottery took place (Perlès and Vitelli, 1999). However, indications of pottery exchange, in the form of imported pottery is found in the Middle Neolithic (MN) where stylistic developments were shared, and small-scale exchanges took place (Perlès and Vitelli, 1999; Pentedeke, 2011; Urem-Kotsou *et al.*, 2012). Analogous traits between vessel technology, shapes and cultural styles infers networks of communication and exchange

operational on regional and inter-regional levels at this time (Çilingiroglu, 2010; Dimoula *et al.*, 2012; Urem-Kotsou *et al.*, 2016). Analysis of lithics from across the period have shown that stone tools were produced in the vicinity, from locally sourced raw materials (Perlès and Vitelli, 1999; Pentedeka, 2011). In the late Neolithic (LN), common pottery wares (black burnished and red slipped) are widespread, thus making it difficult to determine if the pots are locally produced or exchanged (Perlès and Vitelli, 1999). Widespread distribution of materials with limited production sites, such as obsidian, flint and jasper have been identified in the LN (Perlès and Vitelli, 1999; Milić, 2014) as have the movement of stone and spondylus shell ornaments, which circulated over large distances (Pentedeka, 2011; Veropoulidou, 2012). This demonstrates contact between human groups, however, given the small sample size and the number of sites and locations, any suggestion of diachronic change should arguably be made more cautiously. In contrast, $^{87}\text{Sr}/^{86}\text{Sr}$ values from human and animal teeth can be used to infer mobility over longer distances.

Thus far, the use of strontium isotope ratios to identify mobility during the Neolithic of Greece has been restricted to Crete. This small scale study demonstrated that mobility in Crete during the Neolithic was limited or that movements occurred between areas of homogenous underlying geology (Triantaphyllou *et al.*, 2015). To date, no isotopic studies have been conducted on other Neolithic Greek assemblages and aside from movements identified through material culture (Perlès, 2001; Urem-Kotsou *et al.*, 2012; Triantaphyllou *et al.*, 2015) little is known about human mobility in northern Greece. A greater number of mobility studies using strontium isotope ratios have been conducted on Bronze Age tooth enamel and modern reference material from sites situated in the southern Greek mainland and Crete (Nafplioti, 2008; Nafplioti, 2009, 2011). Consequently, this study of strontium isotope analysis on human tooth enamel aims to shed light on the extent of mobility throughout the Neolithic of northern Greece.

The study area in the central and western regions of Greek Macedonia comprises fertile basins separated by mountain ranges. Three broad NE-SW trending geographical ranges give rise to three geological zones bounded by faults which share a common history of deposition and formation (Higgins and Higgins, 1996). The western Pelagonian zone, comprising Triassic and Jurassic limestones and marbles deposited over gneiss. The central Vardar zone, once part of the Tethys Ocean, is made up of Mesozoic deep-water sedimentary and ophiolites, occasionally overlain by limestone and Eocene sediments. The eastern Serbo-Macedonian massif is dominated by metamorphic and plutonic rocks uplifted and faulted by Alpine compressions (Higgins and Higgins, 1996).

Strontium isotope analyses were performed on 36 human tooth enamel samples from seven archaeological sites spanning the Early to Late Neolithic of northern Greece (Fig. 1). The sites in this

study were chosen to chronologically span a large period of the Neolithic and, in addition, cover a range of geographical environments and terrains from coastal to inland locations and fertile basins to mountainous localities (Fig. 1). This allowed temporal study of mobility and comparisons between settlements which are in geographically close proximity with those that are spatially apart. The majority of the human remains studied comprise single burials with minimal grave goods located within the settlements (Triantaphyllou, 2001; Bessios *et al.*, 2003; Triantaphyllou, 2008; Ziota *et al.*, 2009). However, at two of the study sites, Stavroupoli and Toumba Kremastis Koiladas, human burials comprise disarticulated remains and articulated burials are rare (Chondrogianni-Metoki, 1999; Triantaphyllou, 2002, 2004, 2008); at Toumba Kremastis Koiladas many of the dead were cremated (Chondrogianni-Metoki, 2009).

Given the complexity of the regional geology, it is difficult to accurately predict the bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values in the vegetation local to each Neolithic site. For LN Makriyalos, values estimated from modern vegetation samples (of deep-rooted trees and ground vegetation, $n=20$), collected from within 15 km of the archaeological site and determined by Vaiglova *et al.* (in preparation), were used for this purpose. These values are also extended to and combined with the information from geographically and geologically similar sites of Revenia and Paliambela to establish a local $^{87}\text{Sr}/^{86}\text{Sr}$ ratio. For six other sites, bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values were estimated from samples of tooth enamel from archaeological specimens of domestic animals recovered by excavation. Cattle were excluded as they have been shown to be herded over long distances through-out Europe during the Neolithic (Knipper, 2009; Towers *et al.*, 2010; Viner *et al.*, 2010; Sjögren and Price, 2013; Gerling *et al.*, 2017). For this reason, teeth of sheep, goats and pigs were selected where possible for the estimation of locally bioavailable Sr values. There is a possibility that these species can be transhumant; seasonal movement of ovicaprids has detected throughout prehistory (Bocherens *et al.*, 2001; Henton, 2012; Makarewicz *et al.*, 2017; Makarewicz and Pederzani, 2017) however, this is not always the case (Bogaard *et al.*, 2014; Gerling *et al.*, 2015; Marciniak *et al.*, 2017). In the Neolithic of Northern Greece there is no evidence for or against the adoption of transhumance for sheep/goat and we will have to experimentally determine (by way of agreement with either pigs or cows) whether this behaviour is practiced and therefore whether they can for these sites be a local reference.

2. Materials and Methods

Sample preparation, Sr purification and isotope ratio measurement closely follow the procedures outlined in Haak *et al.* (2008), Lewis *et al.* (2014) and Lewis *et al.* (2017). Sample preparation and analyses were conducted in the Bristol Isotope Group, School of Earth Sciences, University of Bristol, UK. The exterior of a tooth was cleaned with a stainless-steel grinding burr and a small piece of enamel (~10 mg) removed from the crown of the tooth using a diamond tipped cutting wheel and any

dentine adhering to the enamel was removed. Enamel samples were then rinsed with ultrapure water (18.2 MΩ) and dried. Super high purity acids were used throughout the strontium chemistry and diluted to the desired concentrations using ultrapure water. Enamel samples were weighed into a clean PFA beaker and dissolved in 7M HNO₃, an aliquot of the sample equivalent to 3 mg of solid enamel was taken for ion exchange chromatography. Strontium was separated from the sample matrix using Sr Spec. resin (Eichrom 50 100 μm particle size; Horwitz *et al.*, 1992). Clean Sr Spec. was loaded on to 100 μL PFA micro columns for Sr separation, samples were loaded in 0.5 mL 3M HNO₃, sample matrix was eluted in 2 mL 3M HNO₃ and strontium was then collected in 1.5 mL ultrapure water.

Strontium isotope analyses were carried out using a ThermoFinnigan Triton Thermal Ionisation Mass Spectrometer (TIMS). Samples were loaded on to rhenium filaments in nitric acid form with 1 μL of tantalum pentachloride (TaCl₅) and 1 μL of 10% phosphoric acid following a modified version of Birck (1986). Triton amplifier gains were determined at the start of each sequence of analyses and all beams were collected on 1011 Ω amplifiers. Filaments were heated automatically, including source tuning, and analysis started once a beam current of 80 pA ⁸⁸Sr was reached. Faraday cups were set to collect ⁸⁴Sr, ⁸⁵Rb, ⁸⁶Sr, ⁸⁷Sr and ⁸⁸Sr as a static analysis and data was collected in 30 blocks of 12 cycles with a 4.194 second integration time per cycle. Samples were corrected for mass fractionation using exponential mass fractionation law and ⁸⁶Sr/⁸⁸Sr of 0.1194 (Nier, 1938; Russell *et al.*, 1978). ⁸⁷Sr beams were corrected for isobaric ⁸⁷Rb using the ⁸⁵Rb/⁸⁷Rb value of 2.59265 (Steiger and Jäger, 1977) which is adjusted for instrumental mass bias. All recorded values were normalised to a known standard NIST SRM 987 ⁸⁷Sr/⁸⁶Sr value of 0.710248 (Avanzinelli *et al.*, 2005). Long term reproducibility of NIST SRM 987 measurements made over the course of this study is 0.71022 ± 0.00005 (2SD; n=24). The reproducibility of a NIST SRM 987 passed through each batch of chemistry is 0.71024 ± 0.00001 (2SD; n=6) and modern marine shell measured with each batch of samples is 0.70916 ± 0.00004 (2SD; n=6).

3. Results and discussion

3.1 Establishing a local range of bioavailable strontium using faunal teeth.

‘Bioavailable strontium’ is the range of local strontium available to plants and animals within an ecosystem, which is delivered to the consumer with an accompanying ⁸⁷Sr/⁸⁶Sr isotope value (Bain and Bacon, 1994). Ecological studies have shown the strontium isotope ratio of terrestrial herbivores broadly reflects that of the surrounding bedrock values (Capo *et al.*, 1998). The local environmental strontium isotope ratio value is made up of terrestrial sources primarily from mineral weathering of surrounding bedrock and atmospheric strontium derived from continental dust and precipitation. At coastal locations sea-spray can also contribute to the local ⁸⁷Sr/⁸⁶Sr ratio values (Miller *et al.*, 1993;

Capo *et al.*, 1998; Vitousek *et al.*, 1999; Whipkey *et al.*, 2000). Human and animal tooth enamel from several sites appear to have values close to that of seawater (0.70918-0.79020; McArthur *et al.*, 2001). A sea-spray affect has been observed at distances of up to 50 m from the coast (Whipkey *et al.*, 2000) and on islands in the Outer Hebrides where biosphere samples reflected seawater strontium isotope ratio rather than the more radiogenic underlying gneiss and granite bedrock (Montgomery and Evans, 2006). As rainwater is derived from seawater, high levels of precipitation ($>c.$ 2000 mm a⁻¹) can result in saturation of the soil and contribute to the bioavailable strontium pool (Evans *et al.*, 2010). The humans and animals from Revenia, Paliambela and Makriyalos situated a few km from the coast at the time of occupation (Pappa and Besios, 1999; Krahtopoulou and Veropoulidou, 2016) each display ⁸⁷Sr/⁸⁶Sr values close to that of seawater. At Stavroupoli the ⁸⁷Sr/⁸⁶Sr values of humans and animals are more radiogenic than that of seawater despite the close proximity of the site to the coast (ca. 3.5 km; Grammenos, 2006). The average rainfall for all settlements would be between 350-500 mm per annum (Panagos *et al.*, 2016), hence, high rainfall means that it is unlikely that the ⁸⁷Sr/⁸⁶Sr values are influenced by soil saturation.

The results presented in Table 1 show that, for the majority of sites, the ⁸⁷Sr/⁸⁶Sr values of Neolithic domestic animals fall within the predicted strontium isotope ratios of the surrounding geology, inferring that animal management most likely occurred at the local level. In fact, the ⁸⁷Sr/⁸⁶Sr values of the faunal samples are far less varied than the values estimated from the bedrock. The results also show there is no difference between the transhumant and non-transhumant species. This demonstrates that the local bioavailable Sr isotope baseline should ideally be determined from small fauna and plants, rather than extrapolating from bedrock as this is likely to underestimate the presence of ‘non-local’ individuals. The exception to this is the site of Stavroupoli, a LN tell site spread over 10 ha (Table 1). Here, the ⁸⁷Sr/⁸⁶Sr values of pig (0.71066), sheep (0.71113) and goat (0.71040), are consistent with the surrounding bedrock, which comprises Holocene coastal and Miocene lacustrine and terrestrial deposits. However, a more radiogenic ⁸⁷Sr/⁸⁶Sr value (0.71255) was determined for the cattle tooth which is consistent with both Cenozoic and Mesozoic metamorphic rocks located to the east of the site (Fig. 1). Mesozoic metamorphic and igneous rocks approximately 3 km east of the settlement, provide a nearby likely candidate for the more radiogenic bioavailable ⁸⁷Sr/⁸⁶Sr value. Although this distance is still very close to the site the more radiogenic Sr ratio observed in the cattle could be either a result of cattle grazing on pastures in an area of more radiogenic ⁸⁷Sr/⁸⁶Sr or animal management strategies such as movement between the coastal flats surrounding Stavroupoli and the slopes of the Mount Chortiatitis to the east (which comprise Cenozoic and Mesozoic metamorphic rocks). The seasonal movement of animals between lowlands and mountainous regions has been observed elsewhere during the Neolithic in Europe (Balasse *et al.*, 2002; Bentley *et al.*, 2003; Bentley and Knipper, 2005; Gerling *et al.*, 2017; Makarewicz, 2017). Although it is difficult to be certain due to the very small sample size, it is also possible that the cow was not raised within the local geologic

area and was perhaps brought to Stavroupoli through exchange. Cattle management strategies throughout the Neolithic have been shown to vary, possibly as a result of adaptation to the local environment. For example, in EN Scandinavia (Gron *et al.*, 2016) and LN Britain (Viner *et al.*, 2010) cattle have been shown to have been driven over long distances. There is also evidence of both local management of livestock (Giblin *et al.*, 2013) and transhumance strategies (Bentley *et al.*, 2003; Stephan *et al.*, 2012).

3.2 Mobility in the Neolithic of northern Greece

The strontium isotope analysis of human tooth enamel undertaken herein suggests that population movement within the Neolithic of northern Greece was highly restricted as almost all populations investigated have $^{87}\text{Sr}/^{86}\text{Sr}$ values consistent with individuals having spent their early life in the local area or moving between areas with similar geology and thus similar strontium isotope ratios. An example of the latter possibility is observed in the $^{87}\text{Sr}/^{86}\text{Sr}$ values of the humans from the sites of Revenia and Makriyalos which are in close geographical proximity and have similar bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$, which could potentially mask movement between them.

During the EN there is little variability in human (average 2SD = 187 ppm; n=10) and faunal (average 2SD = 202 ppm; n=6) strontium isotope values. A similar trend is observed in the MN fauna (average 2SD = 146 ppm; n=4). Towards the LN, however, there is more variation in strontium isotope values for both the fauna (average 2SD = 687 ppm; n=14) and humans (average 2SD = 2607 ppm; n=26) possibly reflecting changes in subsistence strategies and mobility behaviour. For example, the variance apparent in the human strontium isotope values for LN Makriyalos could be a consequence of a larger variability in the faunal samples due to changes in herding strategies or movement of people from an area with a similar underlying geology with differing Sr isotope values.

3.3 Identification of non-local individuals in the sample populations

Strontium isotope analyses of human tooth enamel from the northern Greece Neolithic sites demonstrate that most of the individuals sampled exhibited $^{87}\text{Sr}/^{86}\text{Sr}$ values consistent with having spent their early life in the 'local' area, although movement could have occurred between isotopically homogeneous areas. However, there are a few instances where the $^{87}\text{Sr}/^{86}\text{Sr}$ values of the sample population deviate from the bedrock geology, implying that these individuals spent their early life outside the local area from which their skeletal remains were ultimately recovered.

A notable departure from the general trend of local origin is seen in the population from Kleitos. The site is located on Pleistocene lacustrine and terrestrial sediments. Jurassic and Cretaceous hills sit

close to the north of the site (Fig. 1). The $^{87}\text{Sr}/^{86}\text{Sr}$ values of the local bioavailable Sr estimate at Kleitos are consistent with the upper range of Mesozoic sediments and lowest estimates of Cenozoic sediments and in broad agreement with a sediment source for the Pleistocene age lacustrine and terrestrial deposits (Voerkelius *et al.*, 2010). Consistent with the general trend for Neolithic northern Greece, five of the individuals investigated displayed strontium isotope values that lie within the local $^{87}\text{Sr}/^{86}\text{Sr}$ range determined by the faunal teeth, suggesting that all were resident in the area at the time of tooth mineralisation, or had travelled from an area with a similar bioavailable strontium isotope ratio. Interestingly, however, the strontium isotope ratios of three individuals from Kleitos lie outside of the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ range (Fig. 2). These three individuals are interpreted as having spent their early lives on a more radiogenic geology. The $^{87}\text{Sr}/^{86}\text{Sr}$ values of the possible migrants are consistent with values established for Lower Palaeozoic rocks (Voerkelius *et al.*, 2010). The closest Lower Palaeozoic rocks lie approx. 60 km the north and north-east. The tooth with the most radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ value (0.71467) comes from KL-1, an individual of 6 years of age and of unknown sex. Enamel was sampled from the mandibular premolar, which mineralises between 1.5 and 7 years (Logan and Kronfeld, 1933; Gustafson and Koch, 1974; Schroeder, 1991). This suggests that the child had lived in a different geologic environment for the majority of their young life. Individual KL-2, a male in his mid-30's to mid-40's in age exhibited a slightly less radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ value (0.71285). Enamel was sampled from the maxillary right 2nd molar, which mineralises between 1.5 and 7.5 years (Logan and Kronfeld, 1933; Gustafson and Koch, 1974; Schroeder, 1991). This indicates that the individual was also a non-local during infancy. The final sample, KL-6 is slightly more radiogenic than the local strontium range (0.71089), the individual is estimated to be 15 years of age and of unknown sex. Enamel was sampled from the maxillary left 1st incisor, which mineralises between 3 months and 6 years of age (Logan and Kronfeld, 1933; Gustafson and Koch, 1974; Schroeder, 1991). The interpretation for this individual is more nuanced. It is possible that this individual has had a variable diet with some strontium being sourced from the local area and some from a non-local source either as a direct function of mobility, imported foodstuffs or the individual may have been raised in a region of discrete geology with a bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ of ~ 0.7109 . This value would be consistent with Cenozoic age sedimentary rocks.

The strontium isotope ratios of 5 individuals from Makriyalos lie outside of the determined local isotopic range (Fig. 2), but within the estimated $^{87}\text{Sr}/^{86}\text{Sr}$ values for Miocene age uplands to the west of the site (0.70901 to 0.71100; Voerkelius *et al.*, 2010). Makriyalos, Paliambela and Revenia sit approx. 2-3 km from the same Miocene age upland, so one might expect the biologically available strontium isotope ratios the sites to be similar (~ 0.7092 to 0.7096). This interpretation is supported by the analysis of modern plants used to determine the local bioavailable strontium isotope ratio in this study (0.70900 to 0.70974; Vaiglova *et al.*, in preparation) Teeth MAK-5, 6, 9, 10 and 12 all belong to adults, suggesting movement from an area of a similar geology after tooth mineralisation.

The strontium isotope values for the individuals at Nea Nikomedeia lie outside the estimated $^{87}\text{Sr}/^{86}\text{Sr}$ values for the area for Cenozoic sediments (0.70901 to 0.71100; Voerkelius *et al.*, 2010), but lie in the range predicted for Mesozoic geology (0.70701 to 0.70900), which is consistent with the sediments which form the alluvial plain. The geology surrounding Nea Nikomedeia indicates that the site sits on Pliocene and Quaternary rocks, with alluvial sediment deposits consisting of lacustrine silts and terrestrial clays (Fig. 2). The alluvial plain was formed during the Holocene by the deposition of sediment from four of the surrounding rivers from the Cretaceous and Jurassic uplands to the West (Bintliff, 1976) and thus the source of bioavailable strontium at the site. Although direct determination of local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values from local vegetation would be required to confirm this. Therefore, the individuals at Nea Nikomedeia have $^{87}\text{Sr}/^{86}\text{Sr}$ values consistent with individuals having spent their early life in the local area.

4. Conclusions

In this investigation strontium isotope ratios have been used to explore the movement of Neolithic people in the north of Greece. Baseline bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values calculated from faunal tooth enamel were shown to fall within with predicted $^{87}\text{Sr}/^{86}\text{Sr}$ values of the surrounding bedrock, suggesting that animal management occurred on a local level, although differences in herd management practices became more evident through more varied strontium isotope values at LN sites, which is potentially indicating of changing herding practices leading to a larger variability in the strontium isotope values of the faunal samples. Analyses of human tooth enamel suggests that overall population movement within the region was limited as at almost all sites most individuals exhibit $^{87}\text{Sr}/^{86}\text{Sr}$ values consistent with them having spent their early life in the local area from where their remains were recovered; although the potential for movement between geologically homogenous areas exhibiting similar strontium isotope ratios must always be considered. The strontium isotope ratios of 3 individuals at Kleitos and lie outside of the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ range. These individuals are interpreted as having spent their early lives on a more radiogenic geology, providing evidence of mobility in Greece determined using strontium isotope ratios. The results determined using Sr isotope analysis strengthens the current archaeological evidence for movement and exchange observed through studies of pottery circulation. With both suggesting limited movement in the Early and Middle Neolithic and greater movement in the Late Neolithic.

330 Acknowledgements

331

332 The Natural Environment Research Council (NERC; NE/K500823/1) and the European Union

333 (European Social Fund – ESF) and Greek national funds (NSRF) are thanked for funding the Ph.D.

334 studentship of HLW. HLW would like to thank Dr Thomas Kador for training in strontium isotope

335 analysis.

336

References

- Avanzinelli, R., Boari, E., Conticelli, S., Francalanci, L., Guarnieri, L., Perini, G., Petrone, C., Tommasini, S. & Ulivi, M., 2005. High precision Sr, Nd and Pb isotopic analyses using the new generation Thermal Ionisation Mass Spectrometer ThermoFinnigan Triton-Ti. *Period. Mineral.* 74, 147-166.
- Bain, D.C. & Bacon, J.R., 1994. Strontium isotopes as indicators of mineral weathering in catchments. *Catena*. 22, 201-214.
- Balasse, M., Ambrose, S.H., Smith, A.B. & Price, T.D., 2002. The Seasonal Mobility Model for Prehistoric Herders in the South-western Cape of South Africa Assessed by Isotopic Analysis of Sheep Tooth Enamel. 29, 917-932.
- Balter, V., Braga, J., Telouk, P. & Thackeray, J.F., 2012. Evidence for dietary change but not landscape use in South African early hominins. *Nature*. 489, 558-60.
- Bentley, R.A., Krause, J., Price, T.D. & Kaufman, A., 2003. Human mobility at the Early Neolithic settlement of Vaihingen, Germany. Evidence from strontium isotope analysis. *Archaeometry*. 45, 471-486.
- Bentley, R.A. & Knipper, C., 2005. Geographical Patterns in Biologically Available Strontium, Carbon and Oxygen Isotope Signatures in Prehistoric Sw Germany*. 47, 629-644.
- Bentley, R.A., 2006. Strontium isotopes from the earth to the archaeological skeleton: A review. *J Archaeol Method Th*. 13, 135-187.
- Bentley, R.A., 2013. Mobility and the diversity of Early Neolithic lives: Isotopic evidence from skeletons. *J. Anthropol. Archaeol*. 32, 303-312.
- Bessios, M., Athanassiadou, A., Noulas, K. & Christakou-Tolia, M., 2005. Excavations on the site of the aqueduct in the north of Pieria prefecture (in Greek). *A.E.M.Th.* 17, 2003 Thessaloniki. Aristotle University of Thessaloniki, 451-548.

369

370 Bintliff, J., 1976. The plain of western Macedonia and the Neolithic site of Nea Nikomedeia. Proc
371 Natl Acad Sci U S A. 42, 241-262.

372

373 Bocherens, H., Mashkour, M., Billiou, D., Pellé, E. & Mariotti, A., 2001. A new approach for
374 studying prehistoric herd management in arid areas: intra-tooth isotopic analyses of archaeological
375 caprine from Iran. 332, 67-74.

376

377 Bogaard, A., Henton, E., Evans, J.A., Twiss, K.C., Charles, M.P., Vaiglova, P. & Russell, N., 2014.
378 Locating Land Use at Neolithic Çatalhöyük, Turkey: The Implications of $^{87}\text{Sr}/^{86}\text{Sr}$ Signatures in
379 Plants and Sheep Tooth Sequences. 56, 860-877.

380

381 Boric, D. & Price, T.D., 2013. Strontium isotopes document greater human mobility at the start of the
382 Balkan Neolithic. Proc Natl Acad Sci U S A. 110, 3298-303.

383

384 Britton, K., Grimes, V., Niven, L., Steele, T.E., McPherron, S., Soressi, M., Kelly, T.E., Jaubert, J.,
385 Hublin, J.J. & Richards, M.P., 2011. Strontium isotope evidence for migration in late Pleistocene
386 Rangifer: implications for Neanderthal hunting strategies at the Middle Palaeolithic site of Jonzac,
387 France. 61, 176-85.

388

389 Capo, R.C., Stewart, B.W. & Chadwick, O.A., 1998. Strontium isotopes as tracers of ecosystem
390 processes: theory and methods. Geoderma. 82, 197-225.

391

392 Chondrogianni-Metoki, A., 2001. Excavation on the Prehistoric site of the Kremasti-Kilada Tell,
393 Kozani Prefecture (in Greek). *A.E.M.Th.* 13, 1999 Thessaloniki. Aristotle University of Thessaloniki,
394 399-414.

395

396 Chondrogianni-Metoki, A., 2009. Non residential space uses the Neolithic settlements : the example
397 of Toumba Kremastis Valley (in Greek). PhD thesis, Aristotle University of Thessaloniki (AUTH).

398

399 Çilingiroglu, Ç., 2010. The appearance of impressed pottery in the Neolithic Aegean and its
400 implications for maritime networks in the eastern mediterranean. *Tüba-Ar.* 13.

401

402 Dimoula, A., Pentedeka, A. & Filis, K., 2014. Lete I. The pottery of a neolithic site in central
 403 Macedonia 100 years after. In: Stefani, E., Merousis, N. & Dimoula, A., eds. *A century of research in*
 404 *Prehistoric Macedonia*, 2012 Thessaloniki, Greece. Thessaloniki: Archaeological Museum of
 405 Thessaloniki, 491-503.
 406
 407 Evans, J.A., Montgomery, J., Wildman, G. & Boulton, N., 2010. Spatial variations in biosphere
 408 $^{87}\text{Sr}/^{86}\text{Sr}$ in Britain. *J. Geol. Soc.* 167, 1-4.
 409
 410 Gerling, C., 2015. Prehistoric mobility and diet in the west Eurasian Steppes 3500 to 300 BC. An
 411 isotopic approach, Berlin, De Gruyter.
 412
 413 Gerling, C., Bánffy, E., Dani, J., Köhler, K., Kulcsár, G., Pike, A.W.G., Szeverényi, V. & Heyd, V.,
 414 2015. Immigration and transhumance in the Early Bronze Age Carpathian Basin: the occupants of a
 415 kurgan. 86, 1097-1111.
 416
 417 Gerling, C., Doppler, T., Heyd, V., Knipper, C., Kuhn, T., Lehmann, M.F., Pike, A.W.G. & Schibler,
 418 J., 2017. High-resolution isotopic evidence of specialised cattle herding in the European Neolithic. 12,
 419 e0180164.
 420
 421 Giblin, J.I., Knudson, K.J., Bereczki, Z., Palfi, G. & Pap, I., 2013. Strontium isotope analysis and
 422 human mobility during the Neolithic and Copper Age: a case study from the Great Hungarian Plain. *J.*
 423 *Archaeol. Sci.* 40, 227-239.
 424
 425 Grammenos, D.V., 2006. Conclusions from the rescue excavations at the Neolithic settlement of
 426 Stavroupoli, Thessaloniki. *Analele Banatului*. 14.
 427
 428 Gron, K.J., Montgomery, J., Nielsen, P.O., Nowell, G.M., Peterkin, J.L., Sørensen, L. & Rowley-
 429 Conwy, P., 2016. Strontium isotope evidence of early Funnel Beaker Culture movement of cattle. *J.*
 430 *Archaeol. Sci. Rep.* 6, 248-251.
 431
 432 Grupe, G., Price, T.D., Schröter, P., Söllner, F., Johnson, C.M. & Beard, B.L., 1997. Mobility of Bell
 433 Beaker people revealed by strontium isotope ratios of tooth and bone: a study of southern Bavarian
 434 skeletal remains. *Appl. Geochem.* 12, 517-525.

435
436 Gustafson, G. & Koch, G., 1974. Age estimation up to 16 years of age based on dental development.
437 25, 297-306.

438
439 Haak, W., Brandt, G., de Jong, H.N., Meyer, C., Ganslmeier, R., Heyd, V., Hawkesworth, C., Pike,
440 A.W., Meller, H. & Alt, K.W., 2008. Ancient DNA, strontium isotopes, and osteological analyses
441 shed light on social and kinship organization of the later Stone Age. *Proc Natl Acad Sci U S A.* 105,
442 18226-31.

443
444 Henton, E., 2012. The combined use of oxygen isotopes and microwear in sheep teeth to
445 elucidate seasonal management of domestic herds: the case study of Çatalhöyük, central Anatolia. 39,
446 3264-3276.

447
448 Henton, E., Ruben, I., Palmer, C., Martin, L., Garrard, A., Thirlwall, M. & Jourdan, A.-L., 2017. The
449 Seasonal Mobility of Prehistoric Gazelle Herds in the Azraq Basin, Jordan: Modelling Alternative
450 Strategies Using Stable Isotopes. 23, 187-199.

451
452 Higgins, M.D. & Higgins, R., 1996. A geological companion to Greece and the Aegean, London, UK,
453 Duckworth.

454
455 Hoppe, K.A., Koch, P.L., Carlson, R.W. & Webb, S.D., 1999. Tracking mammoths and mastodons:
456 Reconstruction of migratory behavior using strontium isotope ratios. *Geology.* 27, 439-442.

457
458 Horwitz, P.E., Chiarizia, R. & Dietz, M.L., 1992. A novel strontium-selective extraction
459 chromatographic resin*. *Solvent Extr. Ion Exc.* 10, 313-336.

460
461 Knipper, C., 2009. Mobility in a sedentary society: Insights from isotope analysis of LBK human and
462 animal teeth. In: Hofmann, D. & Bickle, P. (eds.) *Creating Communities: New advances in Central*
463 *European Neolithic Research.* Oxford: Oxbow Books, pp.

464
465 Krahtopoulou, A. & Veropoulidou, R., 2016. Late Pleistocene-Holocene shoreline reconstruction and
466 human exploitation of molluscan resources in northern Pieria, Macedonia, Greece. *J. Archaeol. Sci.*
467 *Rep.*

Lewis, J., Coath, C.D. & Pike, A.W.G., 2014. An improved protocol for $^{87}\text{Sr}/^{86}\text{Sr}$ by laser ablation multi-collector inductively coupled plasma mass spectrometry using oxide reduction and a customised plasma interface. 390, 173-181.

Lewis, J., Pike, A.W.G., Coath, C.D. & Evershed, R.P., 2017. Strontium concentration, radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) and stable ($\delta^{88}\text{Sr}$) strontium isotope systematics in a controlled feeding study. 3, 53-65.

Logan, W.H.G. & Kronfeld, R., 1933. Development of the Human Jaws and Surrounding Structures from Birth to the Age of Fifteen Years**From the Research Department of the Chicago College of Dental Surgery, Dental Department of Loyola University. Read at the Third General Meeting of the Seventy-Fourth Annual Session of the American Dental Association, Buffalo, N. Y., Sept. 14, 1932. 20, 379-428.

Makarewicz, C.A., 2017. Sequential $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ analyses of early Holocene bovid tooth enamel: Resolving vertical transhumance in Neolithic domesticated sheep and goats. 485, 16-29.

Makarewicz, C.A., Arbuckle, B.S. & Öztan, A., 2017. Vertical transhumance of sheep and goats identified by intra-tooth sequential carbon ($\delta^{13}\text{C}$) and oxygen ($\delta^{18}\text{O}$) isotopic analyses: Evidence from Chalcolithic Köşk Höyük, central Turkey. 86, 68-80.

Makarewicz, C.A. & Pederzani, S., 2017. Oxygen ($\delta^{18}\text{O}$) and carbon ($\delta^{13}\text{C}$) isotopic distinction in sequentially sampled tooth enamel of co-localized wild and domesticated caprines: Complications to establishing seasonality and mobility in herbivores. 485, 1-15.

Marciniak, A., Evans, J., Henton, E., Pearson, J., Lisowski, M., Bartkowiak, M. & Sobkowiak-Tabaka, I., 2017. Animal husbandry in the Early and Middle Neolithic settlement at Kopydłowo in the Polish lowlands. A multi-isotope perspective. 9, 1461-1479.

McArthur, J.M., Howarth, R.J. & Bailey, T.R., 2001. Strontium isotope stratigraphy: LOWESS version 3: Best fit to the marine Sr-isotope curve for 0–509 Ma and accompanying look-up table for deriving numerical age. J. Geol. 109, 155-170.

Milić, M., 2014. PXRf characterisation of obsidian from central Anatolia, the Aegean and central Europe. 41, 285-296.

Miller, E.K., Blum, J.D. & Friedland, A.J., 1993. Determination of soil exchangeable-cation loss and weathering rates using Sr isotopes. *Nature*. 362, 438-441.

Montgomery, J. & Evans, J., 2006. Immigrants on the Isle of Lewis - combining traditional funerary and modern isotope evidence to investigate social differentiation, migration and dietary change in the Outer Hebrides of Scotland. In: Gowland, R. & Knusel, C. (eds.) *The Social Archaeology of Funerary Remains*. Oxbow Books, pp. 122-142

Nafplioti, A., 2008. "Mycenaean" political domination of Knossos following the Late Minoan IB destructions on Crete: negative evidence from strontium isotope ratio analysis ($^{87}\text{Sr}/^{86}\text{Sr}$). 35, 2307-2317.

Nafplioti, A., 2009. Mycenae Revisited Part 2. Exploring the Local versus Non-local Geographical Origin of the Individuals from Grave Circle A: Evidence from Strontium Isotope Ratio ($^{87}\text{Sr}/^{86}\text{Sr}$) Analysis. 104, 279-291.

Nafplioti, A., 2011. Tracing population mobility in the Aegean using isotope geochemistry: a first map of local biologically available $^{87}\text{Sr}/^{86}\text{Sr}$ signatures. 38, 1560-1570.

Nier, A.O., 1938. The isotopic constitution of strontium, barium, bismuth, thallium and mercury. *Phys. Rev.* 54, 275-278.

Panagos, P., Ballabio, C., Borrelli, P. & Meusburger, K., 2016. Spatio-temporal analysis of rainfall erosivity and erosivity density in Greece. *Catena*. 137, 161-172.

Pappa, M. & Besios, M., 1999. The Neolithic settlement at Makriyalos, northern Greece: Preliminary report on the 1993-1995 excavations. *J. Field. Archaeol.* 26, 177-195.

532 Pentedeka, A., 2011. Links of clay in Neolithic Greece. The case of Platia Magoula Zarkou. In:
533 Brysbaert, A. (ed.) Tracing Prehistoric social networks through technology: A diachronic perspective
534 on the Aegean. Routledge, pp. 106-125

535

536 Perlès, C. & Vitelli, K., 1999. Craft specialization in the Neolithic of Greece. In: Halstead, P. (ed.)
537 Neolithic Society in Greece. Sheffield: Sheffield Academic Press, pp. 96-107

538

539 Perlès, C., 2001. The Early Neolithic in Greece, Cambridge, UK, Cambridge University Press.

540

541 Russell, W.A., Papanastassiou, D.A. & Tombrello, T.A., 1978. Ca isotope fractionation on the Earth
542 and other solar system materials. *Geochim. Cosmochim. Acta.* 42, 1075-1090.

543

544 Schroeder, H.E., 1991. Oral Structure Biology: Embryology, Structure, and Function of Normal Hard
545 and Soft Tissues of the Oral Cavity and Temporomandibular Joints, G. Thieme Verlag.

546

547 Sjögren, K.G. & Price, T.D., 2013. A complex Neolithic economy: isotope evidence for the
548 circulation of cattle and sheep in the TRB of western Sweden. *J. Archaeol. Sci.* 40, 690-704.

549

550 Steiger, R.H. & Jäger, E., 1977. Subcommittee on geochronology: Convention on the use of decay
551 constants in geo- and cosmochronology. *Earth. Planet. Sci. Lett.* . 36, 359-362.

552

553 Stephan, E., Knipper, C., Schatz, K., Price, T.D. & Hegner, E., 2012. Strontium isotopes in faunal
554 remains: Evidence of the strategies for land use at the Iron Age site Eberdingen-Hochdorf (Baden-
555 Württemberg, Germany). In: Kaiser, E., Burger, J. & Schier, W. (eds.) *Population Dynamics in*
556 *Prehistory and Early History. New Approaches by Using Stable Isotopes and Genetics.* Berlin,
557 Boston: De Gruyter, pp. 265-286

558

559 Towers, J., Montgomery, J., Evans, J., Jay, M. & Pearson, M.P., 2010. An investigation of the origins
560 of cattle and aurochs deposited in the Early Bronze Age barrows at Gayhurst and Irthlingborough. *J.*
561 *Archaeol. Sci.* 37, 508-515.

562

563 Triantaphyllou, S., 2001. A bioarchaeological approach to prehistoric cemetery populations from
564 central and western Greek Macedonia, Oxford, Archaeopress.

565

566 Triantaphyllou, S., 2002. Preliminary analysis of the Neolithic skeletal material from Stavroupoli
567 Thessaloniki (in Greek). In Rescue excavations at the Neolithic settlement of Stavroupolis
568 Thessaloniki. Oxford Archaeopress, pp.

569

570 Triantaphyllou, S., 2004. The Neolithic site of Stavroupoli, excavation season 2002-2003. The study
571 of the human remains (in Greek). In Rescue excavations at the Neolithic Settlement of Stavroupoli,
572 Thessaloniki, part II. pp.

573

574 Triantaphyllou, S., 2008. Living with the dead: a re-consideration of mortuary practices in the Greek
575 Neolithic. In: Isaakidou, V. & Tomkins, P. (eds.) Escaping the labyrinth: The Cretan Neolithic in
576 context. pp. 139-157

577

578 Triantaphyllou, S., Nikita, E. & Kador, T., 2015. Exploring mobility patterns and biological affinities
579 in the Southern Aegean: first insights from Early Bronze Age eastern Crete. The Annual of the British
580 School at Athens. 110, 3-25.

581

582 Urem-Kotsou, D., Papaioannou, A., Papadaku, T., Saridaki, N. & Intze, Z., 2014. Pottery and
583 stylistic boundaries. Early and middle neolithic pottery in Macedonia. In: Stefani, E., Merousis, N. &
584 Dimoula, A., eds. *A century of research in Prehistoric Macedonia*, 2012 Thessaloniki. 505-517.

585

586 Urem-Kotsou, D., Dimoula, A., Elezi, G., Papadaku, T., Papaioannou, A., Saridaki, N., Siamidou, I.,
587 Silva, T., Tzemopoulou, E. & Kotsakis, K., 2016. Patterns in contemporaneous ceramic traditions:
588 inter-regional relations between Thessaly and Macedonia during the Early and Middle Neolithic. In:
589 Sarris, A., Kalogiropoulou, E., Kalayici, T. & Karimali, E. (eds.) *Communities, Landscapes and*
590 *Interaction in Neolithic Greece*. Archaeopress, pp.

591

592 US Geological survey 1:1.5M world geology map, 2017. Available:
593 <http://portal.onegeology.org/OnegeologyGlobal/>.

594

595 Veropoulidou, R., 2014. Molluscan exploitation in the Neolithic and Bronze Age communities at the
596 former Thermaic Gulf, North Aegean. In: Touchais, G., Laffineur, R. & Rougemont, F., eds.
597 *AEAEUM 37: Annales liégeoises et PASPiennes d'archéologie égéenne*, 2012 Paris. 415-422.

598

599 Viner, S., Evans, J., Albarella, U. & Parker Pearson, M., 2010. Cattle mobility in prehistoric Britain:
600 strontium isotope analysis of cattle teeth from Durrington Walls (Wiltshire, Britain). *J. Archaeol. Sci.*
601 37, 2812-2820.

602

603 Vitousek, P.M., Kennedy, M.J., Derry, L.A. & Chadwick, O.A., 1999. Weathering versus atmospheric
604 sources of strontium in ecosystems on young volcanic soils. *Oecologia*. 121, 255-259.

605

606 Voerkelius, S., Lorenz, G.D., Rummel, S., Quetel, C.R., Heiss, G., Baxter, M., Brach-Papa, C.,
607 Deters-Itzelsberger, P., Hoelzl, S., Hoogewerff, J., et al., 2010. Strontium isotopic signatures of
608 natural mineral waters, the reference to a simple geological map and its potential for authentication of
609 food. *Food Chem.* 118, 933-940.

610

611 Whipkey, C.E., Capo, R.C., Chadwick, O.A. & Stewart, B.W., 2000. The importance of sea spray to
612 the cation budget of a coastal Hawaiian soil: a strontium isotope approach. *Chem. Geol.* . 168, 37-48.

613

614 Ziota, C., Chondrogianni-Metoki, V. & Mangouretsiou, E., 2013. The archaeological research in
615 Kleitos Kozani 2009 (in Greek). *A.E.M.Th.* 23, 2009. Aristotle University of Thessaloniki, 37-52.

616

617

618

619 Figure 1
620 Geological map of the study area with the location of settlements. 1. Stavroupoli (LN), 2. Nea
621 Nikomedeia (EN), 3. Paliambela (MN), 4. Makriyalos (LN), 5. Revenia (EN), 6. Kleitos (LN), 7.
622 Toumba Kremastis Koiladas (LN) (base map from US Geological survey 1:1.5M world geology map,
623 2017).

624 Figure 2
625 Scatterplot displaying the variation of human and faunal $^{87}\text{Sr}/^{86}\text{Sr}$ values with the local bioavailable
626 $^{87}\text{Sr}/^{86}\text{Sr}$ ratio through the EN to LN of northern Greece. The shaded area represents the local
627 bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ range defined using strontium isotope ratios of the faunal teeth ($\bar{x} \pm 2\text{SD}$).

628
629
630
631
632

Figure 1





Figure 2



641 Table 1

642 Summary of the $^{87}\text{Sr}/^{86}\text{Sr}$ values for faunal and human teeth by site.

643

644 Table 2

645 Sample information and strontium isotope ratios for human archaeological tooth enamel

646

Site	Faunal $\bar{x} \pm 2SD$	Bioavailable $^{87}Sr/^{86}Sr$ range*	Human $\bar{x} \pm 2SD$	Sediment source rocks	Predicted $^{87}Sr/^{86}Sr$ range***
Nea Nikomedeia	-	-	0.70872 \pm 0.00012	Mesozoic sediments	0.70701 to 0.70900
Revenia	0.70947 \pm 0.00020	0.70927 to 0.70967	0.70944 \pm 0.00024	Pliocene and Quaternary sediments	0.70901 to 0.71100
Paliambela	0.70930 \pm 0.00012	0.70917 to 0.70942	-	Cenozoic sediments	0.70901 to 0.71100
Kleitros	0.70898 \pm 0.00076	0.70822 to 0.70974	0.71046 \pm 0.00444	Cenozoic and Mesozoic sediments	0.70701 to 0.71100
Makriyalos	-	0.70889 to 0.70970**	0.70944 \pm 0.00078	Cenozoic coastal sediments	0.70901 to 0.71100
Stavroupoli	0.71073 \pm 0.00074	0.70999 to 0.71148	0.71113	Cenozoic lacustrine and terrestrial deposits	0.70901 to 0.71100
Toumba Kremastis Koiladas	0.70900 \pm 0.00056	0.70844 to 0.70957	0.70914	Cenozoic sediments and Mesozoic metamorphic rocks	0.70701 to 0.71100

648

649 * in all cases except for Makriyalos, the range of values refers to estimates based on the local
650 bioavailable $^{87}Sr/^{86}Sr$ range defined using strontium isotope ratios of the faunal teeth ($\bar{x} \pm 2SD$).

651 ** range established using measured values of modern vegetation collected within 15 km of the
652 archaeological site (Vaiglova *et al.* in prep).

653 *** Values from Voerkelius *et al.* (2010)

654

Site	Sample	Tooth	Age (yrs)	Sex	$^{87}\text{Sr}/^{86}\text{Sr}$	± 2 SE
Kleitos	KL-1	mandibular P	6.5-7	?	0.71467	0.00001
	KL-2	maxillary RM2	mid 30's-mid 40's	M	0.71285	0.00001
	KL-3	mandibular RP1	late 30's	F	0.70937	0.00001
	KL-4	maxillary RM2	30-40	M	0.70973	0.00001
	KL-5	maxillary RI2	40-50	F	0.70847	0.00001
	KL-6	maxillary LI1	15	?	0.71089	0.00001
	KL-7	mandibular RC	late 30's	F	0.70895	0.00001
	KL-8	maxillary RM2	17-18	M	0.70876	0.00001
Makriyalos	MAK-1	mandibular LM2	-	-	0.70903	0.00001
	MAK-2	mandibular RP2	-	F?	0.70925	0.00001
	MAK-3	mandibular LP2	-	-	0.70917	0.00001
	MAK-4	mandibular RM2	-	-	0.70927	0.00000
	MAK-5	mandibular RM1	18-30	M?	0.70997	0.00001
	MAK-6	mandibular RC	30-40	-	0.71001	0.00001
	MAK-7	mandibular LM2	-	-	0.70930	0.00001
	MAK-8	mandibular RI2	18-30	-	0.70941	0.00001
	MAK-9	maxillary LI1	18-30	-	0.70978	0.00001
	MAK-10	mandibular LP2	18-30	-	0.71054	0.00001
	MAK-11	mandibular RM1	18-30	-	0.70940	0.00001
	MAK-12	maxillary LM1	18-30	-	0.70993	0.00001
	MAK-13	mandibular RM1	-	-	0.70931	0.00001
	MAK-14	mandibular RM1	-	-	0.70964	0.00001
	MAK-15	mandibular RM1	-	-	0.70926	0.00001
	MAK-16	maxillary RM2	-	-	0.70931	0.00001
Nea Nikomedeia	NIK-1	molar	-	-	0.70872	0.00001
	NIK-2	molar	-	-	0.70875	0.00001
	NIK-3	molar	-	-	0.70877	0.00001
	NIK-4	molar	7-8	?	0.70863	0.00001
Revenia	REV-1	mandibular RC	12-13	?	0.70924	0.00001
	REV-2	maxillary LI1	33-35	M	0.70940	0.00001
	REV-3	maxillary LM3	40-50	F?	0.70946	0.00001

	REV-4	mandibular LP2	late 30's	F?	0.70947	0.00001
	REV-5	mandibular LC	40-50	F?	0.70943	0.00012
	REV-6	mandibular LP2	40-50	F?	0.70961	0.00001
Stavroupoli	STAV-5	premolar	-	-	0.71113	0.00001
Toumba						
Kremastis	KK-7	premolar	-	-	0.70914	0.00001
Koiladas						
